


or a pharmaceutically acceptable salt thereof.

19. The compound of claim 1, having the structure



wherein


Ar is a phenyl ring which may be optionally mono-, di- or tri-substituted with a substituent selected from the group consisting of halogen, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, azido, hydroxyalkyl of 1-6 carbon atoms, halomethyl, alkoxymethyl of 2-7 carbon atoms, alkanoyloxymethyl of 2-7 carbon atoms, alkoxy of 1-6 carbon atoms, alkylthio of 1-6 carbon atoms, hydroxy, trifluoromethyl, cyano, nitro, carboxy, alkoxy carbonyl of 2-7 carbon atoms, alkanoyl of 2-7 carbon atoms, benzoyl, amino, alkylamino of 1-6 carbon atoms, dialkylamino of 2-12 carbon atoms, alkanoylamino of 1-6 carbon atoms, alkenoylamino of 3-8 carbon atoms, alkynoylamino of 3-8 carbon atoms, alkanoyloxy of 1-6 carbon atoms, alkenoyloxy of 3-8 carbon atoms, alkynoyloxy of 3-8 carbon atoms, carbamoyl, N-alkylcarbamoyl of 2-7 carbon atoms, N,N-dialkylcarbamoyl of 3-13 carbon atoms, and benzoylamino; or

Ar is the radical:





R₂ is hydrogen, amino, hydroxyamino, trifluoromethyl, alkyl of 1-6 carbon atoms, cycloalkyl of 3-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6



carbon atoms, alkenyloxy of 2-6 carbon atoms, hydroxyalkyl of 1-6 carbon atoms, mercaptoalkyl of 1-6 carbon atoms, halomethyl, alkoxymethyl of 2-7 carbon atoms, alkoxy of 1-6 carbon atoms, cycloalkoxy of 3-8 carbon atoms, alkylthio of 1-6 carbon atoms, cycloalkylthio of 3-8 carbon atoms, alkylsulphinyl of 1-6 carbon atoms, alkylsulfonyl of 1-6 carbon atoms, alkylsulfonamido of 1-6 carbon atoms, alkenylsulfonamido of 2-6 carbon atoms, alkynylsulfonamido of 2-6 carbon atoms, cyano, carboxy, alkoxycarbonyl of 2-7 carbon atoms, alkanoyl of 2-7 carbon atoms, alkanoyloxy of 1-6 carbon atoms, alkenoyloxy of 3-8 carbon atoms, alkynoyloxy of 3-8 carbon atoms, carbamoyl, N-alkylcarbamoyl of 2-7 carbon atoms, N,N-dialkylcarbamoyl of 3-13 carbon atoms, N-alkyl-N-alkenylamino of 4 to 12 carbon atoms, N,N-dialkenylamino of 6-12 carbon atoms, phenylamino, benzylamino, phenoxy, phenyl, thiophenoxy, benzyl, alkylamino of 1-6 carbon atoms, dialkylamino of 2 to 12 carbon atoms, alkanoyloxy of 1-6 carbon atoms, alkenoyloxy of 3-8 carbon atoms, alkynoyloxy of 3-8 carbon atoms, carbamoyl, N-alkylcarbamoyl of 2-7 carbon atoms, N,N-dialkylcarbamoyl of 3-13 carbon atoms,



20



R₃ is hydrogen, alkyl of 1-6 carbon atoms, cycloalkyl of 3-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, hydroxyalkyl of 2-6 carbon atoms; mercaptoalkyl of 2-6 carbon atoms, phenyl, benzyl,



or a pharmaceutically acceptable salt thereof.

10

20. The compound of claim 1, which is:

15

- a) 4-(4-phenoxyanilino)benzo[g]quinoline-3-carbonitrile,
- b) 4-(3-chloro-4-fluoroanilino)benzo[g]quinoline-3-carbonitrile,
- c) 4-(4-chloro-5-methoxy-2-methylanilino)benzo[g]quinoline-3-carbonitrile,
- d) 7,8-dimethoxy-4-(4-phenoxyanilino)benzo[g]quinoline-3-carbonitrile,
- e) 4-(4-chloro-5-methoxy-2-methylanilino)-7,8-dimethoxybenzo[g]quinoline-3-carbonitrile,
- f) 4-(3-chloro-4-fluoroanilino)-7,8-dimethoxybenzo[g]quinoline-3-carbonitrile,
- g) 4-(2,4-dichloroanilino)-7,8-dimethoxybenzo[g]quinoline-3-carbonitrile,
- h) 4-(2,4-dichloroanilino)-7,8-dihydroxybenzo[g]quinoline-3-carbonitrile,

20